



JONATHAN PAYNE, MD

Jonathan Payne, MD

1010 Prince Ave., Suite 700

Athens, GA 30606

Office: (706) 613-5880

---

# What to Expect: Intra-articular Hip Steroid Injection

## What it's for

Most commonly used for:

- **Hip osteoarthritis**
- **Inflammatory arthritis**
- **Diagnostic purposes** (to confirm the hip joint is the pain source)
- Pain limiting walking, sleep, or physical therapy progress

Because the hip joint is deep, **ultrasound guidance** improves accuracy and safety.

---

## Before the injection

- **Medications & conditions to mention:**
  - Diabetes (blood sugar may rise temporarily)
  - Blood thinners
  - Prior hip infection or surgery
- **Clothing:** Wear loose pants or shorts.
- **No fasting** is required.
- You will be asked to **avoid strenuous activity for 48 hours.**

---

## During the procedure (about 10–15 minutes)

1. **Positioning:**
  - You'll lie on your back.
2. **Ultrasound setup:**

- Gel is applied and the hip joint is visualized in real time.
- 3. **Skin cleaning:**
  - Thorough antiseptic prep.
- 4. **Local anesthetic:**
  - You'll feel a brief pinch or burn at the skin.
- 5. **Joint injection:**
  - A longer needle is guided into the hip joint under ultrasound.
  - Sensations may include **pressure, deep ache, or fullness** in the groin or hip.
  - Discomfort is usually brief and tolerable.

Unlike shoulder or knee injections, hip injections can feel **more pressure-like** because of the depth—but ultrasound keeps it quick and precise.

---

#### Immediately after

- **Numbness or heaviness** in the hip or leg from anesthetic (a few hours).
  - Some people feel **immediate pain relief**—this is from the anesthetic, not the steroid.
  - You'll usually be observed briefly, then can go home.
- 

#### First 24–72 hours

- **Soreness at the injection site** or deep hip ache is common.
  - **Steroid flare:**
    - Temporary worsening of pain in ~5–10% of patients.
    - Usually resolves within **48–72 hours**.
  - **Activity:**
    - Light walking is okay.
    - Avoid long walks, running, heavy lifting, or intense exercise for **24–48 hours**.
  - **Ice:**
    - 15–20 minutes at a time if sore.
- 

#### When relief begins

- **Steroid effect:** Usually starts **2–5 days** after injection.
  - **Peak benefit:** Around **1–2 weeks**.
  - **Duration of relief:**
    - Highly variable: **weeks to several months**.
    - Better and longer relief is more common in inflammatory conditions than advanced arthritis.
-

### Common side effects (usually temporary)

- Injection-site pain
- Facial flushing (1–2 days)
- Trouble sleeping the night of injection
- Temporary increase in blood sugar (important for diabetics)

### Less common

- Skin thinning or color change at injection site
- Temporary hip weakness

---

### Rare but serious — seek urgent care if you have:

- Increasing hip pain with **fever**, redness, or warmth (possible infection)
- Severe pain that keeps worsening after 2–3 days
- Inability to bear weight

---

### How often can hip injections be done?

- Usually limited to **2–3 injections per year in the same hip**.
- Repeated injections too frequently may accelerate cartilage wear.
- Dr. Payne recommends **waiting 3 months from the time of the injection until total joint replacement** due to increased risk of infection

---

### How to get the most benefit

- Use the pain-relief window to start or progress **physical therapy**.
- Focus on **hip mobility, glute strengthening, and gait mechanics**.
- Remember: the injection reduces inflammation—it does **not** reverse arthritis or repair structural damage.